LICDAVIS	Coccidioidomycosis Serology Laboratory
CCDAVIO	Serology Laboratory
UNIVERSITY OF CALIFORNIA	Scrutogy Laboratory

Ship specimens frozen or w icepack (49 CFR 173.199 or IATA PI 602 compliant): Lab Phone (530) 752-1757 FedEx, UPS, Coccidioidomycosis Serology Lab, Tupper Hall Room 3137, UC Davis, (530) 752-6813 Fax or Courier West Health Sciences Dr., Davis CA 95616 **Email** coccy@ucdavis.edu Coccidioidomycosis Serology Lab, Tupper Hall Room 3137, UC Davis, USPS www.ucdmc.ucdavis.edu/medmicro/cocci.html 1 Shields Avenue, Davis CA 95616 PATIENT NAME (LAST, FIRST, MI) ORDERING PHYSICIAN (LAST, FIRST, MI) NPI# CLIENT PATIENT ID (MRN) ORDERING PHYSICIAN FAX GENDER DATE OF BIRTH ICD-10 DIAGNOSIS CODE(S) Пм ENTITY TO BE BILLED (IF DIFFERENT THAN ORDERING FACILITY) NO PATIENT BILLING SEND REPORT TO (FACILITY NAME, ADDRESS, STATE, ZIP, EMAIL) SEND REPORT TO (FAX + OPTIONAL EMAIL OR WEB ACCOUNT) E-FAX COMPATIBLE? YES □NO SPECIMEN INFORMATION SPECIMEN ID# COLLECTION DATE SPECIMEN TYPE Serum Other: Cerebrospinal fluid Lumbar Cisternal 3 mL (MINIMUM 2mL) SPECIMEN REQUIRED WITH REQUESTS FOR IMMUNODIFFUSION AND COMPLEMENT FIXATION Ventricular (VP Shunt) TUBES MUST BE LABELED WITH AT LEAST 2 PATIENT IDENTIFIERS Date of Illness Onset SPECIMENS MUST BE SHIPPED IN COMPLIANCE WITH 49 CFR, PART 173.199 OR IATA PACKING INSTRUCTION 602. SERUM STABILITY: 6 MONTHS FROZEN, 2 WEEKS AT 4°C, 48HR AT ROOM TEMPERATURE Patient Race ESR **TOTAL WBC** DIFF: Neut Lymph Mono CSF: Glucose Protein Cells Immunocompromising conditions/drugs Histo Other Coccy serologic results obtained elsewhere: ID Latex EIA:(IgG) EIA:(IgM) Coccidioidal skin test Brief History/Current patient status: Anti-fungal Medications (include total dose to date and route): **TEST(S) REQUESTED** CPT MIN. DAYS **TEST** COST **TEST INFORMATION** TAT CODE(S) VOLUME **PERFORMED** Macrotiter tube-based semi-quantitative IgG assay. In some Coccidioidal complement 2-4 \$44 86171 situations, 'quantitative immunodiffusion' may also be performed at T, W, Th, F fixation (CF) days no additional cost. Coccidioidal CF titers correlate with prognosis. Qualitative IgG and IgM. Specimens are concentrated prior to testing, Coccidioidal immunodiffusion 2-3 \$36 86331 1 ml increasing sensitivity. Coccidioidal immunodiffusion is more sensitive M - F (ID) days and specific than complement fixation. Reflex: Immunodiffusion to \$36 86331 Immunodiffusion is performed and complement fixation is only 2-5 Complement fixation (Default (+ \$44 if 86171 performed (and billed) if immunodiffusion is positive. days if no order selected) reflexed) REQUIRED FOR ALTERNATE (MEDICARE, INSURANCE, PO BILLING) ORDERING FACILITY BILLED IF INCOMPLETE - ATTACHED INSURANCE CARDS OK MEDICARE OR MEDI-CAL# PATIENT ADDRESS INSURANCE PROVIDER PATIENT SOCIAL SECURITY # INSURANCE PROVIDER ADDRESS FOR MINORS OR DEPENDENTS: GUARANTOR NAME & DOB: GROUP # AND SUBSCRIBER ID FOR MINORS OR DEPENDENTS: GUARANTOR ADDRESS PURCHASE ORDER NUMBER (PO #) IF APPLICABLE

NPI#

CLIA#

Director NPI#

1386895068

05D0644125

1831206812